

<https://doi.org/10.15517/rev.biol.trop..v71i1.51382>

Community structure of raptors in the *páramo* landscape of the Ecuadorian Andes

Santiago Barros^{1*};  <http://orcid.org/0000-0003-2647-6579>
Paul Porras¹;  <http://orcid.org/0000-0002-0428-9070>
Boris Landázuri¹;  <http://orcid.org/0000-0001-5034-124X>
David C. Siddons¹;  <https://orcid.org/0000-0003-3305-2969>
Steven C. Latta²; <https://orcid.org/0000-0003-3789-9470>
Pedro X. Astudillo¹;  <https://orcid.org/0000-0002-9945-9414>

1. Laboratorio de Ecología, Escuela de Biología, Universidad del Azuay, Cuenca, Ecuador; jsanty.b1@gmail.com (Correspondence*); paulporraspolo@gmail.com; borisland004@gmail.com; dsiddons@uazuay.edu.ec; pastudillo@uazuay.edu.ec
2. National Aviary, Allegheny Commons West, Pittsburgh, Pennsylvania, USA; steven.latta@aviary.org

Received 01-VII-2022. Corrected 05-X-2022. Accepted 13-IV-2023.

ABSTRACT

Introduction: Habitat alterations result in biodiversity loss, particularly in regions with high levels of diversity and endemism. Raptors are an essential part of the functionality and stability of ecosystems and indicators of habitat quality. In the *paramo* grassland ecosystems in the high Andes of Northern South America, raptors contain a high concentration of threatened species.

Objective: To describe the raptor community structure and determine the species associations.

Methods: We made monthly raptor counts in eight transects from October 2021 to September 2022 and used a principal component analysis to determine species associations.

Results: We identified 149 individuals (seven species, three families) in two communities: abundant (Carunculated Caracara, Variable Hawk, Andean Condor and Turkey Vulture; PCI = 47 %), and scarce (Cinereous Harrier, Peregrine Falcon and Aplomado Falco; PCII = 27 %).

Conclusion: We provide a valid description and understanding of raptor community structure, identifying two communities and the dynamics between them. The first is characterized by an increased abundance of generalist and regionally common species, when the abundance of these species decreases, the second community is defined, characterized by an increase in the abundance of specialist and rare species at the local scale.

Key words: Macizo del Cajas biosphere reserve; protected areas; field methods; raptor census; community analysis.

RESUMEN

Estructura de la comunidad de rapaces en el paisaje de páramo de los Andes ecuatorianos.

Introducción: Las alteraciones del hábitat provocan la pérdida de biodiversidad, especialmente en regiones con altos niveles de diversidad y endemismo. Las aves rapaces son una parte esencial de la funcionalidad y estabilidad de los ecosistemas, y son indicadores de la calidad del hábitat. En los ecosistemas de páramo en los Andes del norte de Sudamérica, hay una concentración de especies rapaces amenazadas.



Objetivo: Describir la estructura de la comunidad de aves rapaces y determinar las asociaciones entre las especies.

Métodos: Hicimos conteos mensuales de rapaces en ocho transectos, de octubre 2021 a setiembre 2022 y usamos un análisis de componentes principales para determinar las asociaciones entre especies.

Resultados: Identificamos 149 individuos (siete especies, tres familias) en dos comunidades: abundantes (e.g., Caracara Curiquingue, Gavilán Variable, Cóndor Andino y Gallinazo Cabecirrojo; PCI = 47 %), y poco abundantes (e.g., Caracara Curiquingue, Gavilán Variable, Cóndor Andino y Gallinazo Cabecirrojo; PCII = 27 %).

Conclusiones: Nuestro enfoque proporciona una descripción y comprensión válida de la estructura de la comunidad de rapaces. Identificamos dos comunidades y la dinámica entre ellas. La primera se caracteriza por una mayor abundancia de especies generalistas y regionalmente comunes, cuando la abundancia de estas especies disminuye, se define la segunda comunidad, caracterizada por un aumento de la abundancia de especies especialistas y raras a escala local.

Palabras clave: reserva de la biosfera Macizo del Cajás; áreas protegidas; métodos de campo; monitoreo de rapaces; análisis de la comunidad.

INTRODUCTION

Habitat alterations associated with global changes negatively influence ecosystems (Hooper et al., 2012). For instance, in tropical montane areas there is an urgent need to monitor biodiversity (Rakotomalala et al., 2021; Roach et al., 2020). For example, in regions of high diversity and endemism, such as the tropical high Andes, rapid land-use change results in an increase in habitat degradation, and this conversion of natural habitats is linked to a biodiversity loss (Myers et al., 2000). Birds are indicators of ecosystem health, being an integral component of habitats and important to ecosystem services (Hudson et al., 2014; Sekercioglu, 2006). Raptors and scavengers provide important services such as prey control and recycling of carcasses, which controls the spread of diseases (Bregman et al., 2014; Sekercioglu, 2006). Raptors are directly related to ecosystem health and are also considered indicator species because they have different degrees of sensitivity as their presence or absences indicates habitat changes (Butet et al., 2022; Pruscini et al., 2016; Sekercioglu, 2006).

The *páramo* is a distinctive ecosystem of the high Andes of Northern South America (Neill, 1999). Here, raptors stand out as a community with a high concentration of threatened species, whose populations are in persistent decline due to illegal hunting and habitat loss (Ballejo et al., 2018; Naveda-Rodríguez

et al., 2016; Plaza & Lambertucci, 2020). Important threatened species in this community include *Vultur gryphus* (Andean Condor), *Falco peregrinus* (Peregrine Falcon), *Falco femoralis* (Aplomado Falcon) and endemics such as *Phalcoboenus carunculatus* (Caracara Curiquingue) (Fjeldsa & Krabbe, 1990; Freile & Restall 2018; Freile et al., 2019; Ridgely & Greenfield, 2001; Stattersfield et al., 1998). Therefore, to improve monitoring efforts for the high Andean raptor community, formal descriptions of field methods and data analysis of the community assemblage is crucial in order to improve management and conservation plans in the region.

In the *páramo* landscape of the Southern Andes of Ecuador, preliminary monitoring efforts have focused on intensive searches and ecological modelling at a species scale (i.e., species-by-species manner), notably include threatened species such as *V. gryphus* (e.g., Astudillo et al., 2011; Astudillo et al., 2016; Naveda et al., 2016). However, no efforts have concentrated on the entire raptor community. Furthermore, the raptor community is characterised by low population densities across large areas (Newton, 1979; Pruscini et al., 2016), resulting in limited efforts to monitor the entire raptor community, as well as complications in data analyses due to low detections, evidencing the need for robust monitoring and analysing protocols. In this context, the present study assesses the high Andean raptor community in

the *páramo* grassland ecosystem of Southern Andes of Ecuador through annual monitoring across transects. Community structure is then explored by means of multivariate analysis (i.e., principal components) to determine the species associations that comprise the community.

MATERIALS AND METHODS

Study area: Our study was carried out the core area of the Macizo del Cajas biosphere reserve in 31 761 ha (3 % of the biosphere reserve) of protected *páramos* in the province of Azuay, Southern Ecuador (Fig. 1). The study boundaries include the Quimsacocha National Recreation Area and buffer zone (i.e., 10 km radius; 3°3'16" S & 79°14' 56" W). The monitoring encompassed around 7 434 ha (i.e., monitoring area influence; Fig. 1). The landscape is characterized by an irregular topography composed of deep, U-shaped valleys, steep

slopes, with rivers on the valley floors and rock mounds at mountain tops (Rodríguez et al., 2014). The elevational range in the study area is 3 430 m.a.s.l. to 3 900 m.a.s.l. The monthly average temperature is 5.4 °C with a maximum of 13.8 °C and a minimum of -0.9 °C. Annual rainfall varies from 1 000 mm to 1 250 mm. The highest precipitation occurs between December and May (Campozano et al., 2016; Ochoa-Sánchez et al., 2018). The vegetation is dominated by *páramo* grassland (90 % of the vegetation cover), an open habitat with herbaceous plants (mainly of the genus *Calamagrostis*) aggregated in tussocks, and associated with shrubby plants of the genera *Chuquiraga*, *Gynoxys*, *Monticalia*, *Valeriana* and *Loricaria* (Baquero et al., 2004; Neill, 1999).

Bird monitoring: Raptor monitoring followed the protocols described by Astudillo et al., (2011) adapted from Ralph et al., (1996) for the Southern Andes of Ecuador. In total, eight




Fig. 1. Map showing the study area and location of eight transects for raptor monitoring in the *páramo* ecosystem, high Andes of Southern Ecuador. The grey shaded polygon represents the Quimsacocha National Recreation Area. The monitoring area influence is 7 434 ha. The secondary map shows the province of Azuay (grey polygon) in Ecuador.



transects of 2 km in length separated by at least 700 m were established. The distance between transects was defined to ensure data independence. All transects were located in wide-open sites to ensure an observation radius of at least 1 km. To avoid double counting, censuses were carried out by two observers who walked each transect at a constant speed and recorded all raptor individuals seen, including those flying overhead. Transects were surveyed only on days with good weather conditions, avoiding periods of rain, fog, and low visibility. Each transect was conducted for three hours once a month from October 2021 to September 2022, with surveys occurring between the hours of 07:00 to 17:00. The total surveying effort was 288 hours of observation. The transects were walked in a random order each month. Taxonomic identification of species follows the classification of the South American Classification Committee (Remsen et al., 2021).

Data analysis: Total abundance was considered as the sum of all individuals per species and per transect (Nur et al., 1999). Community structure was described by principal component analysis (PCA) (Gewers et al., 2021; Huettmann & Diamond, 2001). This technique is one of the most widely used approaches for the analysis and description for biological community data (Vaughan & Ormerod, 2005). The PCA is effective in solving problems associated with different numbers of variables (e.g., species), multicollinearity and small sample sizes (Graham, 2003; Jankowski et al., 2009). Therefore, we used a PCA ordination, based on a correlation matrix of species abundances against sites. The most important components of this ordination were then chosen via the broken-stick method (Jackson, 1993). Sites and species were projected onto a two-dimensional space defined by the chosen components (i.e., biplot), sites closer to each other within the two-dimensional space are considered more similar (Jankowski et al., 2009; Palacio et al., 2020); this graphical visualization of the biological data allows us to make a global

description of the variation in the community data (i.e., species and sites).

We used the loadings of PCA to interpret the model (correlation coefficient between the species abundance and the principal component scores), which utilizes the relative contribution of each variable to each component (Palacio et al., 2020). To define the raptor community as a species set, we considered only loadings with values ≥ 0.25 as a conservative threshold (i.e., 75 % of the data is retained) to summarise each component.

In order to identify the species that integrate each raptor community, we explored an additional graphical option. We created scatterplot with PCI and PCII scores for each site *versus* the abundance of each species per site to visualise the importance of species comprising each community component (i.e., PCI and PCII). Species within each community were identified via establishing cut-off points; these points are limits of a given graph section where changes in the abundance of species can be easily identified (i.e., through an increase or decrease in the abundance of species, and/or the appearance or disappearance of other species) (Ballance et al., 1997; Huettmann & Diamond, 2001). In other words, the scatterplot allows us to visualise a dramatic or evident change in abundance of species associated with PCI and PCII.

RESULTS

In total, 149 individuals were recorded associated with seven species and three families (Table 1). The most abundant species was *P. carunculatus* (51.7 % of records), followed by *Geranoaetus polyosoma* (Variable Hawk) (27.5 % of records) and *Cathartes aura* (Turkey Vulture) (12.7 % of records). The least abundant was *V. gryphus* (Andean Condor) (4 % of records), *F. femoralis* (Aplomado Falcon) (2 % of records), *Circus cinereus* (Cinereous Harrier) (1.7 % of records) and *F. peregrinus* (Peregrine Falcon) (< 1 % of records).

TABLE 1

List of raptor species and their total abundances recorded at eight transects in the páramo grassland ecosystem, high Andes of Southern Ecuador from October 2021 to September 2022.

Order	Family	Common name	Species	Code	Abundance
Cathartiformes	Cathartidae	Turkey Vulture	<i>Cathartes aura</i>	CAAU	19 (mean = 2.38; ± SD = 1.60)
		Andean Condor	<i>Vultur gryphus</i>	VUGR	6 (mean = 0.75; ± SD = 1.16)
Accipitriformes	Accipitridae	Variable Hawk	<i>Geranoaetus polyosoma</i>	GEPO	41 (mean = 5.13; ± SD = 4.22)
		Cinereous Harrier	<i>Circus cinereus</i>	CICI	2 (mean = 0.25; ± SD = 0.46)
Falconiformes	Falconidae	Aplomado Falcon	<i>Falco femoralis</i>	FAFE	3 (mean = 0.38; ± SD = 0.74)
		Peregrine Falcon	<i>Falco peregrinus</i>	FAPE	1 (mean = 0.13; ± SD = 0.35)
		Carunculated Caracara	<i>Phalcoeboenus carunculatus</i>	PHCA	77 (mean = 9.63; ± SD = 9.29)

Community structure: The first three components of our PCA explain 91 % of the total variance in the occurrence of seven raptor species in eight transects in the páramo grassland ecosystem (Table 2). However, through the broken-stick method, only the first two components were retained (74 %). The first component (PCI = 47.4 %) reflects a gradient of change from lower abundance to increased abundance of *P. carunculatus*, *G. polyosoma*, *V. gryphus* and *C. aura*; while the second component (PCII = 27 %) reflects a gradient of change from lower to higher abundance of *C. cinereus* (Cinereous Harrier), *F. peregrinus* and *F. femoralis* (Table 2).

The community ordination (2D solution) showed that PCI is represented by transects composed of more abundant or common species (e.g., *P. carunculatus* and *G. polyosoma*) and are located towards the right side of the biplot (Fig. 2). The PCII axis is represented by

transects that are composed of the less abundant or rare species (e.g., *C. cinereus* and *F. peregrinus*) and are located towards the upper side of the biplot (Fig. 2).

By visually identification of the species composing the raptor community (Fig. 3), cut-off points were determined for each component. In the community represented by the PCI, the abundance of more common species increases at PCA-derived scores greater than 2.86 (Fig. 3A); while in the community represented by the PCII, the abundance of rare species increases at scores greater than 2 (Fig. 3B); these changes show a gradient of variation in the abundance of the raptor community.

DISCUSSION

The patterns of raptor diversity observed during the surveys are consistent with the general patterns reported for the páramos in the

TABLE 2

Loadings of the first three components of the principal component analysis (PCA) for seven raptor species recorded at eight transects in the páramo grassland ecosystem, high Andes of Southern Ecuador

Species	PCI (47.4 %)	PCII (27 %)	PCIII (16 %)
Turkey Vulture (<i>Cathartes aura</i>)	0.421		0.253
Andean Condor (<i>Vultur gryphus</i>)	0.438		-0.413
Variable Hawk (<i>Geranoaetus polyosoma</i>)	0.448	-0.267	
Cinereous Harrier (<i>Circus cinereus</i>)		0.683	
Aplomado Falcon (<i>Falco femoralis</i>)		0.339	-0.736
Peregrine Falcon (<i>Falco peregrinus</i>)	0.326	0.491	0.343
Carunculated Caracara (<i>Phalcoeboenus carunculatus</i>)	0.531		

The first two PCA components were chosen to determine community structure, no loadings < 0.25 are shown (see methods).




Fig. 2. Ordination biplot of the raptor community recorded in eight transects (green circles) in the *páramo* grassland ecosystem, high Andes of Southern Ecuador. The ordination shows the first two components of the principal component analysis (PCA). The arrows describe the loadings of the species that integrate the raptor community. Blue lines represent the vectors associated with species whose greatest contribution is to the first component of the PCA (PCI) and those in red for the second component of the PCA (PCII). For species codes refer to Table 1.

Southern Andes of Ecuador (e.g., Astudillo et al., 2015). Our analysis of the occurrence of seven raptor species in eight transects in the *páramo* grassland ecosystem of the Macizo del Cajas biosphere reserve revealed two distinct raptor communities. Differences in these two communities as revealed by a PCA are related to species abundance. One community is characterized by a higher abundance of generalist as well as regionally common or uncommon species. However, when the abundance of these common species decreases, the raptor community is characterised by an increase in the abundance of specialist and rare species at the local scale. Competition among species for the limited resources in this ecosystem is probably a potential factor that may explain the variations in the raptor community.

The first community is comprised of species that are more typical of the *páramo* landscape, particular across the study area. Species associations including *G. polyosoma* and *P. carunculatus* are expected, as these raptors are common and widely distributed in *páramo* ecosystem (Astudillo et al., 2015; Freile & Restall, 2018). However, our results also show that the common species that integrate the community may also be associated with species such as *V. gryphus* and *C. aura*. The former species is reported as rare in the *páramos* due to its low population densities (Freile & Restall, 2018; Naveda-Rodríguez et al., 2016), while the latter is generally rare in the *páramo* (Astudillo et al., 2015; Olmedo, 2019). On the other hand, *V. gryphus* and *C. aura* are both scavengers and their occurrence has been previously linked to




Fig. 3. Scatterplot of abundance by species and their relationship to scores derived from principal component analysis (PCA) for the raptor community surveyed at eight transects in the *páramo* grassland ecosystem, high Andes of Southern Ecuador. **A.** corresponds to the community described within the first component of the PCA (PCI) and **B.** is the second community described within the second component of the PCA (PCII). Grey arrows show cut-off points: > 2.86 for PCI and > 2 for PCII (see methods). For species codes refer to Table 1.

the presence of more common species, such as *P. carunculatus* (Astudillo et al., 2011; Stucchi & Figueroa, 2010). Peisley et al., (2017) and Sekercioglu, (2006) mention that raptors indirectly lead other species to prey or food sources. In this context, in our study area, especially when *G. polyosoma* and *P. carunculatus* were sighted together, the presence of *V. gryphus* tended to be more frequent. Our findings demonstrate that a community dominated by common species may also be integrated by less common species important for conservation (e.g., *V. gryphus*) and is therefore a characteristic community of the regional *páramo*.

A second community consists of locally rare species such as *C. cinereus*, *F. femoralis* and *F. peregrinus*. These species forage relatively close to *páramo* ground level, flying at

medium altitude and with a more localized distribution (Freile & Restall, 2018; Ridgely & Greenfield, 2001). Within this context, we consider that competition between species of the first community against species of the second community could play a fundamental role in determining the structure of this second raptor community (Ballejo et al., 2018; Donázar et al., 2016; Han et al., 2021), where competitive exclusion of certain species may occur when they share a similar role (Sergio & Hiraldo, 2008; Vrezec & Tome, 2004). For example, *C. cinereus* was observed more frequently when *P. carunculatus* decreased in abundance at transects, perhaps because both species have similar ground-level foraging habits (Fjeldså & Krabbe, 1990; Freile & Restall, 2018; Ridgely & Greenfield, 2001). However, more



studies are required to clarify this relationship. For example, phylogenetic diversity could be employed to understand whether competition is an important factor in species assemblages (e.g., Han et al., 2021).

Our method of characterizing raptor communities through intensive walking transects surveys and principal component analysis proved to be valid for describing and helping to understand *páramo* raptor community structure. The determination of cut-off points in the analysis is appropriate to interpret and identify changes in the raptor community. Huettmann & Diamond, (2001), for example, mention that this approach can be used when species abundances are very low. Therefore, this approach is especially relevant in habitats where raptors have been little studied due to their low densities and wide distribution ranges such as high Andean ecosystems. In addition, we have shown that this approach represents a more comprehensive monitoring protocol that encompasses the overall community assemblage rather than the more common species-by-species assessment, which fails to take into account possible interactions between species. Furthermore, the approximation of principal components and their cut-off points (based on PCA scores and the relationship with their abundances) can be seen as an alternative for understanding the importance of species even when records are scarce. The *páramo* regional application of the methods described in this study could help to determine species associations, its turnover and overlap at a landscape scale. It can also be used as a tool for recognising conservation priorities for threatened species.

Ethical statement: the authors declare that they all agree with this publication and made significant contributions; that there is no conflict of interest of any kind; and that we followed all pertinent ethical and legal procedures and requirements. All financial sources are fully and clearly stated in the acknowledgements section. A signed document has been filed in the journal archives.

ACKNOWLEDGMENTS

We thank the constant support to our research activities of Jacinto Guillén, Raffaella Ansaloni, and Andrés López, from Universidad del Azuay. We thank Eduardo Barnuevo and Bruno Timbe for their field assistance. We also acknowledge Vicente Jaramillo, Soledad Avendaño, Fernando Carrión and Scott Campbell from Dundee Precious Metals, Ecuador (DPMECUADOR S.A.) as well as Mark Thorpe and Jorge Barreno for their continued support to our research. The logistical support came from Hari González and park manager staff of Quimsacocha National Recreation Area, as well as to Empresa Pública de Telecomunicaciones, Agua Potable, Alcantarillado y Saneamiento de Cuenca (ETAPA-EP). This study was funded by Vicerrectorado de Investigaciones, Universidad del Azuay (under grant 2017-0131; 2018-003; 2022-0054), DPMECUADOR S.A., ETAPA-EP (under grant 78010200001) and the Avian Conservation Endowment of the National Aviary (USA).

REFERENCES

- Astudillo, P. X., Siddons, D. C., Barros-Quito, S., Orellana, J. A., & Latta, S. C. (2016). La distribución potencial del cóndor andino. *El Hornero*, 31(2), 89–95.
- Astudillo, P. X., Tinoco, B. A., Graham, C. H., & Latta, S. C. (2011). Assessing methods for estimating minimum population size and monitoring andean condors (*Vultur Gryphus*) in Southern Ecuador. *Ornitología Neotropical*, 22(2), 257–265.
- Astudillo, P. X., Tinoco, B. A., & Siddons, D. C. (2015). The avifauna of Cajas National Park and Mazán Reserve, southern Ecuador, with notes on new records. *Cotinga*, 37(2015), 2–12.
- Ballejo, F., Lambertucci, S. A., Trejo, A., & De Santis, L. J. M. (2018). Trophic niche overlap among scavengers in Patagonia supports the condor-vulture competition hypothesis. *Bird Conservation International*, 28(3), 390–402. <https://doi.org/10.1017/S0959270917000211>
- Ballance, L. T., Pitman, R. L., & Reilly, S. B. (1997). Seabird community structure along a productivity gradient: importance of competition and energetic. *Ecology*, 78(5), 1502–1518.

- Butet, A., Rantier, Y., & Bergerot, B. (2022). Land use changes and raptor population trends: A twelve-year monitoring of two common species in agricultural landscapes of Western France. *Global Ecology and Conservation*, *34*, e02027.
- Baquero, F., Sierra, R., Ordoñez, L., Tipán, M., Espinosa, L., Rivera, M. B., & Soria, P. (2004). *La vegetación de los Andes del Ecuador. Memoria explicativa de los mapas de vegetación: potencial y remanente a escala 1:250.000 y del modelamiento predictivo con especies indicadoras*. EcoCiencia/CESLA/Corporación EcoPar/MAG SIGAGRO/CDC - Jatun Sachal/División Geográfica - IGM.
- Bregman, T. P., Sekercioglu, C. H., & Tobias, J. A. (2014). Global patterns and predictors of bird species responses to forest fragmentation: implications for ecosystem function and conservation. *Biological Conservation*, *169*, 372–383. <https://doi.org/10.1016/j.biocon.2013.11.024>
- Campozano, L., Cèlleri, R., Trachte, K., Bendix, J., & Samaniego, E. (2016). Rainfall and Cloud Dynamics in the Andes: A Southern Ecuador Case Study. *Advances in Meteorology*, *2016*, 1–15. <https://doi.org/10.1155/2016/3192765>
- Donázar, J. A., Cortés-Avizanda, A., Fargallo, J. A., Margalida, A., Moleón, M., Morales-Reyes, Z., Moreno-Opo, R., Pérez-García, J. M., Sánchez-Zapata, J. A., Zuberogoitia, I., & Serrano, D. (2016). Roles of raptors in a changing world: from flagships to providers of key ecosystem services. *Ardeola*, *63*(1), 181–234. <https://doi.org/10.13157/arla.63.1.2016.rp8>
- Fjeldsa, J., & Krabbe, N. (1990). *Birds of the high Andes*. Zoological Museum and Apollo Books.
- Freile, J. F., & Restall, R. L. (2018). *Birds of Ecuador. Helm field guides*. Bloomsbury Publishing Plc.
- Freile, J. F., Santander, T. G., Jiménez-Uzcátegui, G., Carrasco, L., Cisneros-Heredia, D., Guevara, E. A., Sánchez-Nivicela, M., & Tinoco, B. A. (2019). *Lista Roja de las Aves del Ecuador*. Ministerio del Ambiente, Aves y Conservación, Comité Ecuatoriano de Registros Ornitológicos, Fundación Charles Darwin, Universidad del Azuay, Red Aves Ecuador y Universidad San Francisco de Quito.
- Graham, M. H. (2003). Confronting multicollinearity in ecological multiple regression. *Ecology*, *84*(11), 2809–2815.
- Gewers, F. L., Ferreira, G. R., De Arruda, H. F., Silva, F. N., Comin, C. H., Amancio, D. R., & Costa, L. D. F. (2021). Principal component analysis: A natural approach to data exploration. *ACM Computing Surveys*, *54*(4), 70–34. <https://doi.org/10.1145/3447755>
- Hooper, D. U., Adair, E. C., Cardinale, B. J., Byrnes, J. E. K., Hungate, B. A., Matulich, K. L., Gonzalez, A., Duffy, J. E., Gamfeldt, L., & O'Connor, M. I. (2012). A global synthesis reveals biodiversity loss as a major driver of ecosystem change. *Nature*, *486*(7401), 105–108. <https://doi.org/10.1038/nature11118>
- Huettmann, F., & Diamond, A. W. (2001). Using PCA scores to classify species communities: An example for pelagic seabird distribution. *Journal of Applied Statistics*, *28*(7), 843–853. <https://doi.org/10.1080/02664760120074933>
- Hudson, L. N., Newbold, T., Contu, S., Hill, S. L. L., Lysenko, I., De Palma, A., Phillips, H. R. P., Senior, R. A., Bennett, D. J., Booth, H., Choimes, A., Correia, D. L. P., Day, J., Echeverría-Londoño, S., Garon, M., Harrison, M. L. K., Ingram, D. J., Jung, M., Kemp, V., ... Purvis, A. (2014). The PREDICTS database: A global database of how local terrestrial biodiversity responds to human impacts. *Ecology and Evolution*, *4*(24), 4701–4735. <https://doi.org/10.1002/ece3.1303>
- Han, Z., Zhang, L., Jiang, Y., Wang, H., & Jiguet, F. (2021). Local habitat and landscape attributes shape the diversity facets of bird communities in Inner Mongolian grasslands. *Avian Conservation and Ecology*, *16*(1). <https://doi.org/10.5751/ace-01745-160103>
- Jackson, D. A. (1993). Stopping rules in principal components analysis: A comparison of heuristical and statistical approaches. *Ecological Society of America*, *74*(8), 2204–2214.
- Jankowski, J. E., Ciecka, A. L., Meyer, N. Y., & Rabenold, K. N. (2009). Beta diversity along environmental gradients: Implications of habitat specialization in tropical montane landscapes. *Journal of Animal Ecology*, *78*(2), 315–327. <https://doi.org/10.1111/j.1365-2656.2008.01487.x>
- Myers, N., Mittermeier, R. A., Mittermeier, C. G., Da Fonseca, G. A. B., & Kent, J. (2000). Biodiversity hotspots for conservation priorities. *Nature*, *403*(6772), 853–858.
- Neill, D. A. (1999). Vegetación. In P. M. Jørgensen & S. León-Yáñez (Eds.), *Catalogue of the vascular plants of Ecuador* (pp. 13–25). *Monographs in Systematic Botany from the Missouri Botanical Garden* 75.
- Newton, I. (1979). *Population ecology of raptors*. T & AD Poyser.
- Nur, N., Jones, S. L., & Geupel, G. R. (1999). *Statistical Guide to Data Analysis of Avian Monitoring Programs*. Department of the Interior, Fish and Wildlife Service.
- Naveda-Rodríguez, A., Vargas, F. H., Kohn, S., & Zapata-Ríos, G. (2016). Andean Condor (*Vultur gryphus*) in Ecuador: Geographic distribution, population size and extinction risk. *Plos One*, *11*(3), 1–14. <https://doi.org/10.1371/journal.pone.0151827>
- Olmedo, I. (2019). *Cathartes aura*. In J. F. Freire (Ed.), *Aves del Ecuador*. Museo de Zoología, Pontificia



- Universidad Católica del Ecuador. <https://bioweb.bio/faunaweb/avesweb/FichaEspecie/Cathartes%20aura>, acceso Martes, 4 de Enero de 2022.
- Ochoa-Sánchez, A., Crespo, P., & Célleri, R. (2018). Quantification of rainfall interception in the high Andean tussock grasslands. *Ecohydrology*, *11*(3). <https://doi.org/10.1002/eco.1946>
- Palacio, F. X., Apodaca, M. J., & Crisci, J. V. (2020). *Análisis multivariado para datos biológicos: teoría y su aplicación utilizando el lenguaje R*. Vazques Mazzini Editores.
- Plaza, P. I., & Lambertucci, S. A. (2020). Ecology and conservation of a rare species: What do we know and what may we do to preserve Andean condors? *Biological Conservation*, *251*(2020), 108782. <https://doi.org/10.1016/j.biocon.2020.108782>
- Pruscini, F., Morelli, F., Perna, P., Mazzeo, R., Cavitolto, P., Bertellotti, M., Catorci, A., & Santolini, R. (2016). Roadside raptor surveys in valdes Peninsula (Patagonia, Argentina). *Hornero*, *31*(1), 7–11.
- Peisley, R. K., Saunders, M. E., Robinson, W. A., & Luck, G. W. (2017). The role of avian scavengers in the breakdown of carcasses in pastoral landscapes. *Emu Austral Ornithology*, *117*(1), 68–77. <https://doi.org/10.1080/01584197.2016.1271990>
- Remsen, J. V., Areta, J. I., Bonaccorso, E., Claramunt, S., Jaramillo, A., Lane, D., Pacheco, J., Robbins, M., Stiles, F., & Zimmer, K. (2021). *Una clasificación de las especies de aves de América del Sur*. Unión Americana de Ornitólogos.
- Ridgely, R. S., & Greenfield, P. J. (2001). *The Birds of Ecuador, Volume 1: Status, Distribution, and Taxonomy*. Cornell University Press.
- Ralph, C. J., Geupel, G. R., Pyle, P., Martin, T. E., DeSante, D. F., & Borja, M. (1996). *Manual de métodos de campo para el monitoreo de aves terrestres*. Pacific Southwest Research Station, Forest Service, United States Department of Agriculture Forest.
- Rodríguez, S., Rodas, F., Schubert, A., & Vasco, S. (2014). *Área de biosfera macizo del Cajas; Experiencias de desarrollo sostenible para el buen vivir*. ETAPA EP. Municipio de Cuenca, Ministerio del ambiente, SENPLADES, Ministerio de relaciones exteriores, Cooperación alemana GIZ, Naturaleza y cultura internacional.
- Roach, N. S., Urbina-Cardona, N., & Lacher, T. E. (2020). Land cover drives amphibian diversity across steep elevational gradients in an isolated neotropical mountain range: Implications for community conservation. *Global Ecology and Conservation*, *22*, e00968. <https://doi.org/10.1016/j.gecco.2020.e00968>
- Rakotomalala, A. A. N. A., Wurz, A., Grass, I., Martin, D. A., Osen, K., Schwab, D., Soazafy, M. R., Tschamtké, T., & Raveloson-Ravaomanarivo, L. H. (2021). Tropical land use drives endemic versus exotic ant communities in a global biodiversity hotspot. *Biodiversity and Conservation*, *30*(14), 4417–4434. <https://doi.org/10.1007/s10531-021-02314-4>
- Sekercioglu, C. H. (2006). Increasing awareness of avian ecological function. *Trends in Ecology and Evolution*, *21*(8), 464–471. <https://doi.org/10.1016/j.tree.2006.05.007>
- Stucchi, M., & Figueroa, J. (2010). Descripción de las interacciones tróficas entre el cóndor andino *Vultur gryphus* y otras especies por el consumo de carroña en el cañón del Colca, Arequipa. *Boletín Informativo de la Unión de Ornitólogos de Perú*, *5*(2), 8-14.
- Sergio, F., & Hiraldo, F. (2008). Intraguild predation in raptor assemblages: A review. *Ibis*, *150*(1), 132–145. <https://doi.org/10.1111/j.1474-919X.2008.00786.x>
- Vaughan, I. P., & Ormerod, S. J. (2005). Increasing the value of principal components analysis for simplifying ecological data: A case study with rivers and river birds. *Journal of Applied Ecology*, *42*(3), 487–497. <https://doi.org/10.1111/j.1365-2664.2005.01038.x>
- Vrezec, A., & Tome, D. (2004). Altitudinal segregation between Ural Owl *Strix uralensis* and Tawny Owl *S. aluco*: Evidence for competitive exclusion in raptorial birds: Capsule The Owls were significantly segregated in space with the most important factor being altitude. *Bird Study*, *51*(3), 264–269. <https://doi.org/10.1080/00063650409461362>