

The invasion of five alien species in the Delta do Parnaíba Environmental Protection Area, Northeastern Brazil

Daniel Loebmann¹, Ana Cecília G. Mai² & James T. Lee³

1. Laboratório de Herpetologia, Programa de Pós-Graduação em Ciências Biológicas (Zoologia), Instituto de Biociências, Universidade Estadual Paulista, Av. 24 A, 1515, Bairro Bela Vista, CEP 13506-900, Rio Claro, SP, Brazil; contato@danielloebmann.com
2. Embrapa Meio-Norte, Laboratório de Recursos Aquáticos, BR 343, km 35, CEP 64200-970, Parnaíba, PI, Brazil; anacecilia_mai@yahoo.com.br
3. Laboratório de Biologia Pesqueira e Manejo de Recursos Aquáticos, Universidade Federal do Pará, Av. Perimetral, 2651. CEP 66077-830, Belém, PA, Brazil; james.jtlee@gmail.com

Received 20-IX-2009. Corrected 26-II-2010. Accepted 25-III-2010.

Abstract: Marine biological invasions have been regarded as one of the major causes of native biodiversity loss, with shipping and aquaculture being the leading contributors for the introductions of alien species in aquatic ecosystems. In the present study, five aquatic alien species (one mollusk, three crustaceans and one fish species) were detected during dives, shore searches and from the fisheries on the coast of the Delta do Parnaíba Environmental Protection Area, in the States of Piauí and Maranhão, Northeastern Brazil. The species were the bicolor purse-oyster *Isognomon bicolor*, the whiteleg shrimp *Litopenaeus vannamei*, the giant river prawn *Macrobrachium rosenbergii*, the Indo-Pacific swimming crab *Charybdis hellerii* and, the muzzled blenny *Omobranchus punctatus*. Ballast water (*I. bicolor*, *C. hellerii*, and *O. punctatus*) and aquaculture activities (*L. vannamei* and *M. rosenbergii*) in adjacent areas are the most likely vectors of introduction. All exotic species found have potential impact risks to the environment because they are able to compete against native species for resources (food and habitat). *Isognomon bicolor* share the same habitat and food items with the native bivalve species of mussels and barnacles. *Litopenaeus vannamei* share the same habitat and food items with the native penaeids such as the pinkspot shrimp *Farfantepenaeus brasiliensis*, the Southern brown shrimp *Farfantepenaeus subtilis*, and the Southern white shrimp *Litopenaeus schmitti*, and in the past few years *L. vannamei* was responsible for a viral epidemics in the cultivation tanks that could be transmitted to native penaeid shrimps. *Charybdis hellerii* is also able to cause impacts on the local fisheries as the species can decrease the populations of native portunid crabs which are commercialized in the studied region. *Macrobrachium rosenbergii* may be sharing natural resources with the Amazon River prawn *Macrobrachium amazonicum*. *Omobranchus punctatus* shares habit with the native redlip blenny *Ophioblennius atlanticus* and other fishes, such as the frillfin goby *Bathigobius soporator*. Some immediate remedial measures to prevent further introductions from ballast water and shrimp farm ponds should be: (i) to prevent the release of ballast water by ship/vessels in the region; (ii) to reroute all effluent waters from shrimp rearing facilities through an underground or above-ground dry well; (iii) to install adequate sand and gravel filter which will allow passage of water but not livestock; (iv) outdoor shrimp ponds located on floodable land should be diked, and; (v) to promote environmental awareness of those directly involved with ballast water (crews of ship/vessels) and shrimp farms in the region. Rev. Biol. Trop. 58 (3): 909-923. Epub 2010 September 01.

Key words: exotic species, ballast water, aquaculture, Mollusca, Decapoda, Blenniidae.

Numerous factors have been appointed to explain the disappearance of wild species from their natural habitats. Among them, biological invasions are a leading cause of

species extinctions and biotic homogenization in terrestrial and freshwater systems worldwide (Sala *et al.* 2000, Lockwood & McKinney 2001, but see Gurevitch & Padilla 2004

for marine habitats), ranking second only, to habitat loss as the major threat to biodiversity (Wilson 1992, Wilcove *et al.* 1998).

The magnitude of the problem is extensive, and invasions frequently interact with other factors such as habitat loss, pollution and climate change compromising the integrity of marine ecosystems. Despite the severity of the implications, studies demonstrating the impacts of introduced species in the marine environment are rare, and most introductions probably go unnoticed (Rilov & Crooks 2009). Other factors have also been suggested as facilitators of the establishment of invasive species, including high reproductive rates, facilitated dispersal, low levels of competition among resident species, tolerance to a wide range of environmental conditions, often, and lacking predators in their new environment (Crawley 1986, Kolar & Lodge 2001, Sakai *et al.* 2001).

Several studies have summarized the impacts of invasive species on native species, and their community structure (Williamson 1996, Wilcove *et al.* 1998, Parker *et al.* 1999, Sala *et al.* 2000, Stein *et al.* 2000). These studies attributed the main impacts on native biota to the spread of exotic pathogens and parasites, biological competition for space and food, predation upon native species, and alteration of the genetic makeup in closely related species. Moreover, many species listed as threatened or endangered are considered to be at risk primarily because of competition with and predation by non-indigenous species (Wilcove *et al.* 1998).

Marine and freshwater ecosystems are particularly vulnerable to invasions of alien species as the dispersion of propagules in water occurs easier than in terrestrial environments. Minchin *et al.* (2009) provide a wide review of the vectors that contribute to the invasions of aquatic alien species and point out eight main pathways for a non-native organism to arrive in a new location. Among these vectors, shipping and aquaculture have been considered as the most critical pathways for marine invasions globally (Molnar *et al.* 2008). Burke *et al.* (2000) estimated that the marine ecosystems in the Mediterranean contain 480 invasive

species, the Baltic 89, and Australian waters 124. Considering that the number of invasive species is increasing quickly and that undisturbed natural areas are declining, biological impacts by alien species may become the leading factor of ecological disintegration (Crooks & Soulé 1996).

In Brazil, despite the researchers efforts to try to understand the patterns of dispersion, causes and consequences of bioinvasions, the surveys are very limited. According to Ferreira *et al.* (2009) there is a clear trend of increasing bioinvasion events in regional coastal ecosystems, but whether invasion rates are actually increasing or are a result of more intensive research efforts in recent years is still an open question. Therefore, to understand and to track the expansion of the bioinvasions in the Brazilian coast, we present new records of occurrence of five aquatic alien species in the Environmental Protection Area and propose guidelines to circumvent further introductions to the coastal zone of the States of Piauí and Maranhão (Northeast Brazil), an ecological sanctuary which supports a rich diversity of aquatic fauna, including commercial and threatened species.

MATERIAL AND METHODS

Site characteristics: Established on August 28th 1996, the Delta do Parnaíba Environmental Protection Area (Delta do Parnaíba, EPA) is located among the pairs of geographical coordinates 02°37'-03°05' S and 42°29'-41°09' W. The area covers 3 138km², which comprises the entire littoral of the State of Piauí and parts of the littorals of the States of Ceará and Maranhão. The tidal variation in the area is classified as mesotidal (Davies 1964), with a maximum range of 3.3m. Water temperature is homogenous throughout the area and averages 28.5±0.9°C (mean±SD). According to Köppen's climate classification updated (Kottek *et al.* 2006), the area is classified as Aw (Equatorial climate with dry winter). Temperatures are warm throughout the year ranging from 22 to 33°C with a marked

dry period, when precipitation is lower than 100mm between June and December, and a rainy period in the first months of the year (January-May).

The coastal zone of the Delta do Parnaíba EPA is formed by a complex mosaic of several habitats such as coastal dunes, rocky shores, mangrove forests, estuaries, and sandy beaches. Three estuaries (Parnaíba River Delta, Camurupim/Cardoso and Timonha/Ubatuba) have important roles in the ecology of the coastal communities due to the release of large amounts of sediments and organic matter, especially in the rainy season. In addition, the Parnaíba River Delta is considered the third largest oceanic deltaic formation worldwide.

Sampling methods: Specimens of the non-native species were recorded from February 2005 to April 2009 in the municipalities of Parnaíba, Luis Correia, Cajueiro da Praia (State of Piauí) and Araisos (State of Maranhão). Individuals were obtained by the following methods: (1) Snorkeling; this method was performed sporadically on the beaches of Barra Grande (Cajueiro da Praia) and Coqueiro (Luis Correia) during the low tide (ca. 2 hours each dive); (2) Intensive search for unfamiliar individuals on rocky outcrops, which was conducted monthly on the beaches of Cajueiro da Praia and Barra Grande (Cajueiro da Praia), and Coqueiro (Luis Correia) during the low tide (ca. 3 hours each search); and (3) Monitoring the local fisheries (casting nets) during the season fishery of the white shrimp (May to July). Total length of each specimen collected was measured. Voucher specimens were fixed in 10% formalin, preserved in 70% alcohol, and deposited at the Zoological collection of the Natural History Museum of the Universidade Federal da Bahia (UFBA) and the Crustacean collection of the Fundação Universidade Federal do Rio Grande (FURG). Collecting permits were authorized by the Brazilian Environmental Protection Agency, the Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis (IBAMA) (Proc. Number 18027-2).

RESULTS

Five aquatic alien species (one mollusk, three crustaceans and one fish) new to the study area were identified during the period of this study (Fig. 1), i.e. the bicolor purse-oyster *Isognomon bicolor* (C. B. Adams 1845), the whiteleg shrimp *Litopenaeus vannamei* (Boone, 1931), the giant river prawn *Macrobrachium rosenbergii* (De Man, 1879), the Indo-Pacific swimming crab *Charybdis hellerii* (A. Milne-Edwards, 1867) and, the muzzled blenny *Omobranchus punctatus* (Valenciennes 1836).

Mollusca, Bivalvia, Isognomonidae:

Isognomon bicolor. In June 2008, the mats of *I. bicolor* (Fig. 2) were identified attached on granite rock shores at the Pedra do Sal beach, municipality of Parnaíba, State of Piauí (02°48'10" S, 41°43'46" W) (Table 1). Pedra do Sal is the only beach on the coast of Piauí with granitic rocks permanently exposed. In addition, the beach is typically reflective with high-energy waves acting on the outcrops. The individuals of *I. bicolor* were found on the same substrates of the native mussels (*Mytilus* sp.), in small patches (each <2 m²) at the same zone of *Mytilus* sp.

Crustacea, Decapoda, Dendrobranchiata, Penaeidae: *Litopenaeus vannamei*. During April and May 2009, four specimens of *L. vannamei* (Fig. 3) were collected. The individuals were obtained from casting nets of fishermen in the municipalities of Luis Correia (02°53' S, 41°37' W) and Cajueiro da Praia (02°55' S, 41°21' W) (Fig. 1, Table 2). The individuals of *L. vannamei* were found along with two native shrimp species, the Atlantic seabob *Xiphopenaeus kroyeri* (C. Heller, 1862) and the Southern white shrimp *Litopenaeus schmitti* (Burkenroad, 1936). Individuals collected were adults ranging from 10 to 12.6cm standard length.

Crustacea, Decapoda, Pleocyemata, Carididae: *Macrobrachium rosenbergii*. In April




Fig. 1. Delta do Parnaíba Environmental Protection Area showing the location where the alien specimens were captured. In the map of Brazil, the States with coastal zone are in gray color. From North towards South the States in gray are Pará, Amapá, Maranhão, Piauí, Ceará, Rio Grande do Norte, Paraíba, Pernambuco, Alagoas, Sergipe, Bahia, Espírito Santo, Rio de Janeiro, São Paulo, Paraná, Santa Catarina and Rio Grande do Sul.

2009 three adult males of *M. rosenbergii* were detected in Delta do Parnaíba EPA: one specimen collected by a fisherman using a casting net at Barra Grande Beach, in the municipality of Cajueiro da Praia, State of Piauí (02°54'06" S, 41°24'25" W) (Fig. 4) (Voucher number: UFBA 260; 26.8cm standard length), one specimen captured by fishermen in Tatus' pier harbor (02°49'56" S, 41°49'48" W), Ilha Grande de Santa Isabel municipality, Delta do Parnaíba estuary, and a specimen captured by fishermen in the Carnaubearas, Araisos municipality, border between the States of Piauí and Maranhão (02°49'38" S, 41°57'45" W) (Fig. 1, Table 3).

Crustacea, Decapoda, Pleocyemata, Portunidae: *Charybdis hellerii*. Three specimens (two males and one female) were collected in the municipality of Cajueiro da Praia, State



Fig. 2. *Isognomon bicolor*: Specimen collected at Pedra do Sal beach, State of Piauí, Brazil.

of Piauí (Fig. 5). All specimens were found occupying rock outcrops during the low water (02°54'10" S, 41°24'21" W) (Fig. 1, Table 4). Vouchers specimens were deposited in the

TABLE 1

Isognomon bicolor. *Known records, localities and new records (present study) in Brazil. States: PI-Piauí; RN-Rio Grande do Norte; PE-Pernambuco; BA-Bahia; RJ-Rio de Janeiro; SP-São Paulo; SC-Santa Catarina*

State	Municipality	Latitude (s)	Longitude (w)	Reference
PI	Parnaíba	02°48'10"	41°43'46"	New record
RN	Natal	05°52'	35°09'	1
PE	Recife	08°06'	34°53'	1
BA	Salvador	12°57'	38°21'	1
RJ	Arraial do Cabo	22°57'	42°01'	1
RJ	Niterói	22°58'	43°03'	1
RJ	Ilha Grande	23°05'-23°13'	44°05'-44°22'	2
SP	Ubatuba	23°22'-23°36'	44°43'-45°19'	1
SP	Ilha Bela	23°43'-23°58'	45°13'-45°27'	1
SP	Itanhaém	24°22'	46°48'	3
SP	São Sebastião	23°46'-23°49'	45°26'-45°39'	1
SP	Bertioga	23°47'	45°59'	1
SP	Guarujá	23°58'-24°03'	46°10'-46°15'	1
SP	São Vicente	23°58'	46°22'	1
SC	Itapema	27°05'	48°36'	1
SC	Porto Belo	27°07'	48°31'	1
SC	Bombinhas	27°08'-23°49'	45°26'-45°39'	1
SC	Florianópolis	27°37'	48°26'	1

1. Domaneschi & Martins 2002; 2. Oliveira & Creed 2008; 3. Jacobucci *et al.* 2006.

TABLE 2

Litopenaeus vannamei. *Known records, localities and new records (present study) in Brazil. States: PI-Piauí; RN-Rio Grande do Norte; SP-São Paulo*

State	Municipality	Latitude (s)	Longitude (w)	Reference
PI	Luis Correia	02°53'	41°37'	New records
PI	Cajueiro da Praia	02°55'	41°21'	New records
RN	Canguaretama	06°22'	35°00'	1
RN	Nísia Floresta	06°07'	35°11'	1
RN	Sen. Georgino Avelino	06°09'	35°07'	1
RN	Arêz	06°12'	35°09'	1
SP	Cananéia	24°57'-25°03'	47°48'-47°56'	2

1. Santos & Coelho 2002; 2. Barbieri & Melo 2005.

Crustacean collection (3208) of the Fundação Universidade Federal de Rio Grande.

Chordata, Osteichthyes, Blenniidae:
Omobranchus punctatus. In December 2008, two specimens of *O. punctatus* were collected under the rocks in the intertidal zone at Coqueiro beach (02°53'55" S, 41°34'17" W) (Fig. 6, Table 5), municipality of Luis Correia

(Fig. 1). These specimens were deposited in the Bahia Federal University Zoological Collection (UFBA 5268). The specimens collected measured 8.2 and 9.0cm standard length.

DISCUSSION

Early detection of invasive species is fundamental to develop management actions to




Fig. 3. Specimen of *Litopenaeus vannamei* with standard length=12.6cm (UFBA 259), collected at Barra Grande beach, State of Piauí, Brazil.




Fig. 4. Specimen of *Macrobrachium rosenbergii* with standard length=26.8cm (UFBA 260), collected in Barra Grande beach, municipality of Cajueiro da Praia, State of Piauí, Brazil.

TABLE 3

Macrobrachium rosenbergii. Known records, localities and new records (present study) in Brazil. States: PA-Pará; MA-Maranhão; PI-Piauí; ES-Espírito Santo; SP-São Paulo; PR-Paraná

State	Municipality	Latitude (s)	Longitude (w)	Reference
PA	Colares	00°56'	48°50'	1
PA	Salvaterra	00°45'	48°30'	1
MA	Araioses	02°49'38"	41°57'45"	New records
PI	Ilha Grande	02°49'56"	41°49'48"	New records
PI	Cajueiro da Praia	02°54'06"	41°24'24"	New records
ES	locality not specifield			2
SP	Brejo Alegre	21°07'	50°10'	3
PR	Pontal do Paraná	25°41'	48° 31'	4

1. Barros & Silva 1997; 2. Valent & New 2000; 3. Magalhaes *et al.* 2005; 4. Gazola-Silva *et al.* 2007.

TABLE 4

Charybdis hellerii. Known records, localities and new records (present study) in Brazil. States: MA-Maranhão; PI-Piauí; CE-Ceará; RN-Rio Grande do Norte; PE-Pernambuco; AL-Alagoas; BA-Bahia; RJ-Rio de Janeiro; SP-São Paulo; PR-Paraná; SC-Santa Catarina

State	Municipality	Latitude (s)	Longitude (w)	Reference
MA	São Luis, Raposa	02°27'-02°30'	44°02'-44°17'	1
PI	Cajueiro da Praia	02°54'10"	41°24'21"	New records
PI	Luis Correia	02°54'-02°55'	41°27'-41°33'	2
CE	Icapuí	04°42'	37°21'	3
CE	Fortim	04°27'	37°04'	3
RN	Macau	05°06'-05°07'	36°15'-36°44'	4
PE	Tamandaré	08°44'	35°05'	5
AL	Maceió	locality not specifield		6
BA	Salvador	23°10'	44°30'	7
BA	Camamu	13°56'	39°05'	8
BA	Ilhéus	14°47'	39°01'	9
RJ	Rio de Janeiro	22°46'	43°06'	10
RJ	Angra dos Reis	23°02'	44°21'	10
RJ	Niterói	22°52'	43°06'	10
SP	Ubatuba	23°22'-23°36'	44°43'-45°19'	11
SP	Caraguatatuba	23°37'	45°24'	11
SP	São Vicente	23°58'	46°23'	12
PR	Guaratuba	25°50'	48°34'	13
SC	Florianópolis	27°48'	48°38'	14

1. Feres *et al.* 2007; 2. Lima-Junior *et al.* 2008; 3. Bezerra & Almeida 2005; 4. Ferreira *et al.* 2001; 5. Coelho & Santos 2003; 6. Calado 1996; 7. Carqueija & Gouvêa 1996; 8. Almeida *et al.* 2003; 9. Almeida *et al.* 2006; 10. Tavares & Mendonça, 1996; 11. Braga *et al.* 2005; 12. Reigada *et al.* 2006; 13. Frigotto & Serafim-Junior 2007; 14. Mantelatto & Dias 1999.



Fig. 5. Specimen of *Charybdis hellerii* with carapace width=8.0cm (FURG 3208), collected in a tide pool at Barra Grande beach, State of Piauí, Brazil.



Fig. 6. Specimen of *Omobranchus punctatus* with standard length=8.2cm (UFBA 5268), collected in a tidepool at Coqueiro beach, State of Piauí, Brazil.

avoid the establishment or to control the spreading of invasive populations. This is particularly important because at early time, measures for contention or eradication are most effective. In the present study, five new occurrences of introduced species, varying from three different phyla, gives an indication of the ecological risks the Delta do Parnaíba EPA currently faces. Accordingly, some aspects of the biology

of these species and considerations about their distribution in the Delta do Parnaíba EPA are presented, which could form a baseline for monitoring the impacts of these alien species.

***Isognomon bicolor*:** According to Ferreira *et al.* (2009) *I. bicolor* is one of the most important species introduced to the benthic realm along the Brazilian coast. Originally from the

TABLE 5

Omobranchus punctatus. Known records, localities and new records (present study) in Brazil. States: PI-Piauí; BA-Bahia; RJ-Rio de Janeiro; SC-Santa Catarina

State	Municipality	Latitude (s)	Longitude (w)	Reference
PI	Luis Correia	2°53'55"	41°34'17"	New record
BA	Salvador	23°10'	44°30'	1
RJ	Angra dos Reis	22°50'-23°20'	44°00'-44°45'	2
SC	São Francisco do Sul	26°00'-26°40'	48°28'-49°00'	2

1. Mendonça *et al.* 2005; 2. Gerhardinger *et al.* 2006.

Caribbean region, *I. bicolor* has been detected in Brazilian waters since 1994 (Domaneschi & Martins 2002), i.e. in the states of Rio Grande do Norte, Pernambuco, Bahia, Rio de Janeiro, São Paulo, and Santa Catarina (Domaneschi & Martins *op cit.*, Jacobucci *et al.* 2006, Oliveira & Creed 2008) (Table 1). Ballast water and fouling of commercial vessels are the suggested vectors of introduction (Rocha 2002). Another hypothesis is that *I. bicolor* could have been introduced into Brazil accidentally on oil or gas platforms from the Caribbean coast (Oliveira & Creed 2008).

At the moment, the impact of this species on native biota in the region seems to be low. However, the species has dominated the mid-intertidal rocky shores along the Southeastern coast of Brazil (Magalhães 1999, Rapagnã 2004, Breves-Ramos 2004). For example, López (2003) estimated that a population of the native barnacle *Tetraclita stalactifera* reduced ca. 50% since the invasion of *I. bicolor* in Cabo Frio, State of Rio de Janeiro. Besides the competition and displacement of barnacle populations, *I. bicolor* probably competes with a commercial species of Mytilidae, *Perna perna*, a very abundant mussel found in the region (Rapagnã 2004, Breves-Ramos 2004, Ferreira *et al.* 2009), which is also an exotic species from Africa that reached the Brazilian coast in the 18th and 19th centuries during the slave trade (Ferreira *et al.* *op cit.*).

***Litopenaeus vannamei*:** During the rainy season the probability of shrimp escapes increases considerably because the cultivation

tanks, especially those near the rivers, cannot always support the excessive rainfalls. Fig. 7 shows two shrimp farms installed in the surroundings of Camurupim River where some tanks are within 30m of the river. Consequently, *L. vannamei* could be accidentally released from overflowed ponds and lakes into the marine areas.



Fig. 7. Satellite image showing the proximity of the shrimp farms (*Litopenaeus vannamei* cultivation tanks) at the surroundings of Camurupim River (shrimp ponds less than thirty meters of Permanent Preservation Area). 1- Camurupim river; 2- Cultivate shrimp ponds; and 3- Mangrove forests. Image from Google Earth version 5.0.11733.9347.

Litopenaeus vannamei was imported to Brazil in the 1970's for aquaculture purposes. The species was considered the most viable for cultivation compared to native species due to the high adaptability to the varying hydro-chemical conditions of the tropical estuaries in Northeast Brazil. At the moment, *L. vannamei* has been reported in Brazilian estuarine and

coastal waters from Rio Grande do Norte (Santos & Coelho 2002) to São Paulo (Barbieri & Melo 2005) (Table 2). However, the occurrence of this species in the wild may be larger than documented as it is relatively difficult to distinguish *L. vannamei* from native penaeid species such as *Farfantepenaeus paulensis*, *Farfantepenaeus brasiliensis* and *Farfantepenaeus subtilis*. To what extent these fugitive specimens of poorly managed shrimp farms will be able to establish self-sustaining populations in the natural environment is still an open question (Ferreira *et al.* 2009).

***Macrobrachium rosenbergii*:** The giant river prawn *M. rosenbergii* is the most cultivated freshwater shrimp in the world. It is native to the Indo-Pacific region (Pakistan, Vietnam, Philippines, New Guinea and Australia) (Chan 1998). *Macrobrachium rosenbergii* was brought to Brazil in 1977 for experimental cultivation purposes by the Department of Oceanography of the Universidade Federal de Pernambuco (Cavalcanti 1998). The species disseminated quickly to the States of Maranhão, Ceará, Pernambuco, Rio de Janeiro, São Paulo, and Santa Catarina (Barros & Silva 1997, Cavalcanti 1998, Valenti & New 2000).

Although *Macrobrachium rosenbergii* lives almost the entire life cycle in fresh water, it is considered an estuarine-depend species since it depends on brackish water to reproduce. According to Ling (1969), the physiological dependence of the larvae to brackish water is an important factor to contribute to the establishment of this species in natural waters. In the Brazilian coast there appears to be no well-established population of *M. rosenbergii*, however, records of occurrence in the wild has been observed for the States of Espírito Santo (Valenti & New 2000), São Paulo (Magalhães *et al.* 2005), Paraná (Gazola-Silva *et al.* 2007) and Pará (Barros & Silva 1997) (Table 3). The rainy season of 2009 accumulated the highest levels of rain fall since 1974 in coastal zone of Piauí. Considering that this species is cultivated in aquaculture farms around the City of Teresina (05°05'21" S, 43°48'09" W), it is

probable that the incidental introduction was made by escapees from these farms.

***Charybdis hellerii*:** The portunid crab *Charybdis hellerii* is a voracious Indo-Pacific species. The first documented invasion of *C. hellerii* occurred in Mediterranean Sea after the opening of the Suez Channel in the late 19th century (Galil 1992, Rodriguez & Suárez 2001). In Brazil, the species has been reported since 1995 (Tavares & Mendonça 1996) and thus far, the species has been reported from Maranhão (North) to Santa Catarina (South) (Carqueija & Gouvêa 1996, Calado 1996, Negreiros-Fransozo 1996, Mantelato & Dias 1999; Tavares & Mendonça 1996, Carqueija 2000, Ferreira *et al.* 2001, Almeida *et al.* 2003, Coelho & Santos 2003, Bezerra & Almeida 2005; Braga *et al.* 2005, Almeida *et al.* 2006, Reigada *et al.* 2006, Feres *et al.* 2007, Frigotto & Serafim-Junior 2007, Coelho *et al.* 2008, Lima-Junior *et al.* 2008) (Table 4). According to Ferreira *et al.* (2009) two hypotheses should be considered for the presence of *C. hellerii* in Brazilian waters. First, the possibility that the species arrived in Brazil as a result of natural dispersion from populations inhabiting the Caribbean Seas since the mid-1980s. Second, the invasion of Brazilian waters through secondary introduction with ballast water.

Furthermore, *Charybdis hellerii* might have economic impacts to local fisheries as this species may become a competitor of the native portunid crabs (*Callinectes* spp.) which are commercialized in the region. Although *C. hellerii* is similar in size with the native portunids, this exotic species has no commercial value (Lima-Junior *et al.* 2008).

***Omobranchus punctatus*:** The original distribution of the muzzled blenny *O. punctatus* was presumed to be the Indo-Pacific region, ranging from Japan and Australia to the Persian Gulf (Golani 2004). In Brazil, the first occurrences were registered in the States of Bahia, Rio de Janeiro and Santa Catarina (Mendonça *et al.* 2005, Gerhardinger *et al.* 2006) (Table 5).

Omobranchus punctatus arrived into Brazilian waters by ballast or bilge water, hidden inside the fouling of ships (Gerhardinger *et al.* 2006). However, records presented here may have been an effect of the species dispersion since the maritime transport in the region is restricted to small vessels with limited use of ballast water.

Conservation implications: The introduction of exotic and allochthonous species to local communities and ecosystems has been leading to relevant modifications in the food webs, steady-state populations and changes in the ecosystems functional processes (Rocha *et al.* 2005).

Delta do Parnaíba EPA is clearly vulnerable to aquatic bioinvasions. Two probable sources, aquaculture and shipping, are responsible for the introduction of alien species in the Delta do Parnaíba EPA. *Litopenaeus vannamei* and *Macrobrachium rosenbergii* are species which likely escaped from surrounding aquaculture farms (Santos & Coelho 2002, Barbieri & Melo 2005) and *Isognomon bicolor*, *Charybdis hellerii* and *Omobranchus punctatus* were species likely introduced with ships (Tavares & Mendonça 1996, Gerhardinger *et al.* 2006).

Environmental impacts associated with the implementation of non-sustainable practices in shrimp farms have been neglected in favor to immediate socio-economic benefits. Mangrove deforestation for the establishment of cultivation tanks, outflow of highly eutrophic effluents into natural water bodies, and the potential dissemination of shrimp virus diseases to native crustacean populations are among the main impacts caused by aquaculture (Boeger *et al.* 2005, Ferreira *et al.* 2009).

Tavares & Mendonça (2004) have reported the introduction of at least 18 species of exotic decapods in the Brazilian coastline. Five species (28%) have managed to keep reproductively active populations, *Charybdis hellerii*, *Rhithropanopeus harrisi* (Gould, 1841), *Pyromaia tuberculata* (Lockington, 1877), *Macrobrachium rosenbergii* (De Man, 1879), and *Procambarus clarkii* (Girard, 1852). Two

species have probably reproduced in Brazilian waters, *Pilummoides perlatus* (Poeppig, 1836) and *Penaeus monodon* (Fabricius, 1798); and the remaining 11 species (61%) had few individuals recorded in natural environments.

All exotic species recorded in this study have high potential to compete for natural resources against native fauna. *Isognomon bicolor* share the same habitat and food items with the native species of mussels and barnacles (López 2003, Fernandes *et al.* 2004). *Litopenaeus vannamei* share the same habitat and food items with the native penaeids such as the pinkspot shrimp *Farfantepenaeus brasiliensis* (Latreille, 1817), the Southern brown shrimp *Farfantepenaeus subtilis* (Pérez Farfante, 1967), and the Southern white shrimp *Litopenaeus schmitti* (Burkenroad, 1936). *Charybdis hellerii* share resources (space and food) on rocky shores with several brachyurans such as the masked swimming crab *Callinectes larvatus* Ordway, 1863, the Rugose swimming crab *Callinectes exasperatus* (Gerstaecker, 1856), the Dana swimming crab *Callinectes danae*, the Cuban stone crab *Menippe nodifrons* (Stimpson, 1859), and the narrowback mud crab *Panopeus americanus* Saussure, 1857. *Macrobrachium rosenbergii* may be sharing natural resources with the Amazon River prawn *Macrobrachium amazonicum* (Heller, 1862). *Omobranchus punctatus* shares habit with the native redblip blenny *Ophioblennius atlanticus* (Valenciennes, 1836) and other fishes, such as the frillfin goby *Bathigobius soporator* (Valenciennes, 1837).

Litopenaeus vannamei brought other threats to native crustacean species. In the past few years, almost all farms with cultivated shrimps in the State of Piauí and adjacent states were suffering from a high incidence of viral epidemics, such as the White spot syndrome (WSS), the Taura Syndrome Virus (TSV), the Infectious Myonecrosis Virus (IMNV) and the Necrotizing Hepatopancreatitis (NHP) (Pantoja & Lightner 2008). These pathogens are highly lethal to cultivated *Litopenaeus vannamei* (Briggs *et al.* 2005) and could be transmitted to native penaeid shrimp populations

(Overstreet *et al.* 1997, Briggs *et al.* 2005). Many species have been reported as susceptible hosts (*Penaeus merguensis* De Man, 1888 and *Metapenaeus monoceros* (Fabricius, 1798), while *Palaemon setiferus* Olivier, 1811, *Euphausia superba* Dana, 1852, *Metapenaeus dobsoni* (Miers, 1878), *Parapenaeopsis stylifera* (H. Milne Edwards, 1837), *Solenocera indica* Nataraj, 1945, *Squilla mantis* L., 1758, *Macrobrachium rosenbergii* and other crustaceans can act as latent carriers (Flegel *et al.* 1997, Hossain *et al.* 2001).

The release of non-native organisms is a federal offense according to Brazilian legislation and ballast water/aquaculture in Brazil should be conducted strictly in accordance with specific regulations, control and effective inspections (Alves *et al.* 2007). Some immediate remedial measures to prevent further introductions from ballast water and shrimp farm ponds should be: (i) to prevent the release of ballast water by ship/vessels in the region; (ii) to reroute all effluent waters from shrimp rearing facilities through an underground or above-ground dry well; (iii) to install adequate sand and gravel filter which will allow passage of water but not livestock; (iv) outdoor shrimp ponds located on floodable land should be diked, and; (v) to promote environmental awareness of those directly involved with ballast water (crews of ship/vessels) and shrimp farms in the region.

Otherwise, the bicolor purple-oyster, the whiteleg shrimp, the giant river prawn, the Indo-Pacific swimming crab, and the muzzled blenny escapes out of most of the vessels and shrimp ponds will negatively change the native diversity scenario in the future.

In order to understand the real impacts that alien species may cause after introduction to new environments, biological and ecological studies are still necessary. Therefore, the data presented in this study is of notable importance to environmental institutions of the government (Instituto Brasileiro do Meio Ambiente e Recursos Naturais Renováveis and Instituto Chico Mendes de Conservação da

Biodiversidade) which are the managers of the Delta do Parnaíba EPA.

ACKNOWLEDGMENTS

The authors are grateful to Silmara Erthal (Instituto Chico Mendes de Conservação da Biodiversidade) for providing the shapefile of the Delta do Parnaíba EPA. Claudio Sampaio (Universidade Federal da Bahia) and Cleverton R. M. dos Santos (Museu Paraense Emílio Goeldi) for confirming the identifications of some specimens. Maria Cristina Oddone for preparing the spanish abstract of the manuscript. Conselho Nacional de Pesquisa e Desenvolvimento (CNPq) for providing a doctoral scholarship to Daniel Loebmann (grant no. 140226/2006-0).

RESUMEN

En el Área de Protección Ambiental del Delta Del Río Parnaíba, noreste de Brasil fueron detectadas cinco especies acuáticas invasoras (un molusco, tres crustáceos y un pez): la ostra de dos colores *Isognomon bicolor*, el camarón patiblanco *Litopenaeus vannamei*, el camarón gigante de río *Macrobrachium rosenbergii*, el cangrejo nadador del Indo-Pacífico *Charybdis hellerii* y el blenio hocicudo *Omobranchius punctatus*. El agua de lastre (*I. bicolor*, *C. hellerii*, y *O. punctatus*) y la acuicultura (*L. vannamei* y *M. rosenbergii*) en áreas adyacentes son los vectores de introducción más probables. Todas las especies exóticas encontradas son potencialmente riesgosas para el ambiente ya que son aptas para competir con las nativas por recursos como alimento y hábitat. *Charybdis hellerii* también podría causar impacto en el recurso pesquero local, al reducir las poblaciones de cangrejos portunidos nativos, comercializados en el área de estudio.

Palabras clave: especies exóticas, agua de lastre, acuicultura, Mollusca, Decapoda, Blenniidae.

REFERENCES

- Almeida, A.O., P.A. Coelho & J.T.A. Santos. 2003. New records of decapod crustaceans (Dendrobranchiata and Brachyura) for the state of Bahia, Brazil. *Nauplius* 11: 129-133.
- Almeida, A.O., P.A. Coelho, J.T.A. Santos & N.R. Ferraz. 2006. Crustáceos decápodos estuarinos de Ilhéus, Bahia, Brasil. *Biota Neotrop.* 6: 1-24.

- Alves, C.B.M., F. Vieira, A.L.B. Magalhães & M.F.G. Brito. 2007. Impacts of non-native fish species in Minas Gerais, Brazil: present situation and prospects, p. 291-314. *In* T.M. Bert (ed.). Ecological and genetic implications of aquaculture activities. Springer, Dordrecht, Zuid-Holland, Netherland.
- Barbieri, E. & G.A.S. Melo. 2005. Biodiversity: *Litopenaeus vannamei* (Boone, 1931) exotic species presence in Cananéia-Iguape-Ilha Comprida lagoon estuary complex. *O Mundo da Saúde* 30: 654-659.
- Barros, M.P. & L.M.A. Silva. 1997. Registro de introdução da espécie exótica *Macrobrachium rosenbergii* (De Man, 1879) (Crustacea, Decapoda, Palaemonidae), em águas do estado do Pará, Brasil. *B. Mus. Paraense Emilio Goeldi, série Zoologia* 13: 31-37.
- Bezerra, L.E.A. & A.O. Almeida. 2005. Primeiro registro da espécie Indo-Pacífica *Charybdis hellerii* (A. Milne - Edwards, 1867) (Crustacea: Decapoda: Portunidae) para o litoral do Estado do Ceará. *Trop. Oceanog.* 33: 33-38.
- Boeger, W.A., M. Pie, A. Ostrensky & L.P. Azambuja. 2005. Identity of the putative agent of the lethargic crab disease (LCD), responsible by massive mortalities of the mangrove crab, *Ucides cordatus* (Brachyura, Ocypodidae), in Northeastern Brazil. *Mem. Inst. Oswaldo Cruz* 100: 161-167.
- Braga, A.A., A. Fransozo, G. Bertini & P.B. Fumis. 2005. Composition and abundance of the crabs (Decapoda, Brachyura) off Ubatuba and Caraguatatuba, northern coast of São Paulo, Brazil. *Biota Neotrop.* 5: 1-34.
- Breves-Ramos A. 2004. Distribuição, abundância e estrutura populacional de *Isognomon bicolor* (Adams, 1845) no litoral do estado do Rio de Janeiro, Brasil. MSc. Thesis, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Rio de Janeiro, Brazil.
- Briggs, M., S. Funge-Smith, R.P. Subasinghe & M. Phillips. 2005. Introductions and movement of two penaeid shrimp species in Asia and the Pacific. FAO Fisheries Technical Paper, no. 476, Food and Agriculture Organization of the United Nations, Rome, Lazio, Italy.
- Burke, L., Y. Kura, K. Kassem, M. Spalding & C. Revenga. 2000. Pilot Analysis of Global Ecosystems: Coastal Ecosystems Technical Report. World Resources Institute, Washington, Columbia, USA.
- Calado, T.C.S. 1996. Registro de *Charybdis hellerii* (Milne Edwards, 1867) em águas do litoral brasileiro (Decapoda: Portunidae). *B. Est. Ciênc. Mar* 9: 175-180.
- Carqueija, C.R.G. & E.P. Gouvêa. 1996. A ocorrência, na costa brasileira, de um Portunidae (Crustacea, Decapoda), originário do Indo-Pacífico e Mediterrâneo. *Nauplius* 4: 105-112.
- Cavalcanti, L.B. 1998. Histórico, p. 17-20. *In* W.C. Valenti (ed.). Carcinocultura de Água Doce. Tecnologia para Produção de Camarões. IBAMA/FAPESP, Brasília, Distrito Federal, Brazil.
- Chan, T.Y. 1998. Shrimps and prawns, p. 851-966. *In* K.E. Carpenter & V.H. Niem (eds.). FAO species identification guide for fishery purposes. The living marine resources of the Western Central Pacific. Volume 2. Cephalopods, crustaceans, holoturians and sharks. Food and Agriculture Organization of the United Nations, Rome, Lazio, Italy.
- Coelho, P.A., A.O. De Almeida & L.E.A. Bezerra. 2008. Checklist of the marine and estuarine Brachyura (Crustacea: Decapoda) of Northern and Northeastern Brazil. *Zootaxa* 1956: 1-58.
- Coelho, P.A. & M.C.F. Santos. 2003. Ocorrência de *Charybdis hellerii* (Milne Edwards, 1867) (Crustacea, Decapoda, Portunidae) no litoral de Pernambuco. *Bol. Téc. Cient. CEPENE* 11: 167-173.
- Crawley, M.J. 1986. The population biology of invaders. *Phil. Trans. R. Soc. B.* 314: 711-731.
- Crooks, J. & M.E. Soulé. 1996. Lag times in population explosions of invasive species: Causes and implications, p. 103-125. *In* O.T. Sandlund, P.J. Schei & Å. Viken. The Trondheim Conferences on Biodiversity. Invasive Species and Biodiversity Management. Kluwer Academic, Trondheim, Sør-Trøndelag, Norway.
- Davies, J.L. 1964. A morphogenic approach to world's shorelines. *Zeits Fur Geomorph.* 8: 127-142.
- Domaneschi, C. & C.M. Martins. 2002. *Isognomon bicolor* (C.B. Adams) (Bivalvia, Isognomonidae): primeiro registro para o Brasil, redescricao da espécie e considerações sobre a ocorrência e distribuição de *Isognomon* na costa brasileira. *Rev. Bras. Zool.* 19: 601-610.
- Feres, S.J.C., A.T.L. Lopes & L.A. Santos. 2007. Primeiro registro de *Charybdis hellerii* (Milne Edwards, 1867) para o litoral Maranhense-Brasil (Crustacea, Decapoda, Portunidae). *Bol. Lab. Hidrobiol.* 20: 77-82.
- Fernandes, F.C., L.C. Rapaganã & G.B.D. Bueno. 2004. Estudo da população do bivalve exótico *Isognomon bicolor* (C. B. Adams 1845) (Bivalvia, Isonomoniidae) na Ponta da Fortaleza em Arraial do Cabo-RJ, p. 134-141. *In* J. Silva & R. Souza (eds.). Água de

- Lastro e Bioinvasão. Interciência, Rio de Janeiro, Brazil.
- Ferreira, C.E.L., A.O.R. Junqueira, M.C. Villac & R.M. Lopes. 2009. Marine Bioinvasions in the Brazilian Coast: brief report on history of events, vectors, ecology, impacts and management of non indigenous species, p. 459-478. *In* G. Rilov & J.A. Crooks. Biological invasions of marine ecosystems: ecological, management, and geographic perspectives. Springer-Verlag, Berlin, Brandenburg, Germany.
- Ferreira, A.C., C. Sankarankutty, I.M.C. Cunha & F.T. Duarte. 2001. Yet another record of *Charybdis helleri* (A.Milne Edwards) (Crustacea, Decapoda) from the Northeast of Brazil. *Rev. Bras. de Zool.* 18: 357-358.
- Flegel, T.W., S. Boonyaratpalin & B. Withyachumnarnkul. 1997. Progress in research on yellow-head virus and white-spot virus in Thailand, p. 285-295. *In* T.W. Flegel & I.H. MacRae (eds.). Diseases in Asian Aquaculture III. Fish Health Section. Asian Fisheries Society, Manila, Metro Manila, Philippines.
- Frigotto, S.F. & M. Serafim-Junior. 2007. Primeiro registro de *Charybdis hellerii* (Milne Edwards, 1867) (Crustacea) no litoral do estado do Paraná. *Est. Biol.* 29: 227-230.
- Galil, B. 1992. Eritrean decapods in the Levant. Biogeography in motion. *B. l'Institut. Océanogr.* 9: 115-123.
- Gazola-Silva, F.F., S.G. Melo & J.R.S. Vitule. 2007. *Macrobrachium rosenbergii* (Decapoda: Palaemonidae): possible introduction in a costal river of Paraná (Brazil). *Acta Biol. Parana* 36: 83-90.
- Gerhardinger, L.C., M.O. Freitas, A.B. Andrade & C.A. Rangel. 2006. *Omobranchus punctatus* (Teleostei: Blenniidae), an exotic blenny in the Southwestern Atlantic. *Biol. Inv.* 8: 941-946.
- Golani, D. 2004. First record of the muzzled blenny (Osteichthyes: Blenniidae: *Omobranchus punctatus*) from the Mediterranean, with remarks on ship-mediated fish introduction. *J. Mar. Biol. Ass. U.K.* 84: 851-852.
- Gurevitch, J. & D.K. Padilla. 2004. Are invasive species a major cause of extinctions? *Trend. Ecol. Evol.* 19: 470-474.
- Hossain, M.D.S., A. Chakraborty, B. Joseph, S.K. Otta & I. Karunasagar. 2001. Detection of new hosts for white spot syndrome virus using nested polymerase chain reaction. *Aquaculture* 198: 1-11.
- Jacobucci, G.B., A.Z. Güth, A. Turra, C.A. Magalhães, M.R. Denadai, A.M.R. Chaves & E.C.F. Souza. 2006. Levantamento da macrofauna associada a *Sargassum* spp. na ilha da Queimada Pequena, Estação Ecológica dos Tupiniquins, litoral sul do Estado de São Paulo, Brasil. *Biota Neotrop.* 6: 1-8.
- Kolar, C.S. & D.M. Lodge. 2001. Progress in invasion biology: predicting invaders. *Trend. Ecol. Evol.* 16: 199-204.
- Kottek, M., J. Grieser, C. Beck, B. Rudolf & F. Rubel. 2006. World Map of the Köppen-Geiger climate classification updated. *Meteorol. Z.* 15: 259-263.
- Lima-Junior, T.B., M.I.C. Aragão, J.P. Silva, G.A.S. Melo & J.R.S.A. Leite. 2008. Occurrence of two Indo-Pacific species of Brachyura on the coast of Piauí, Brazil. *Bol. Lab. Hidrob.* 21: 35-40.
- Ling, S.W. 1969. The general biology and development of *Macrobrachium rosenbergii* (De Man). *FAO Fish. Rep.* 57: 589-606.
- López, M.S. 2003. Efecto de la potencial presa exótica *Isognomon bicolor* (Adams, 1845) sobre la ecología trófica de *Stramonita haemastoma* (Kool, 1987) en el intermareal rocoso de Arraial do Cabo, RJ, Brasil. M.Sc. Thesis, Universidad Internacional de Andalucía, Córdoba, Spain.
- Lockwood, J.L. & M.L. McKinney. 2001. Biotic homogenization. *Kluwer, Dordrecht, Zuid-Holland, Netherland.*
- Magalhães, C.A. 1999. Partilha de recursos em guilda de gastrópodes predadores em costões de São Sebastião, SP. Ph.D. Thesis, Universidade Estadual de Campinas, Campinas, São Paulo, Brazil.
- Magalhães, C., S.L.S. Bueno, G. Bond-Bunkup, W.C. Valenti, H.L.M. Silva, F. Kiyohara, E.C. Mossolin & S.S. Rocha. 2005. Exotic species of freshwater decapod crustaceans in the state of São Paulo, Brazil: records and possible causes of their introduction. *Biodiv. Conserv.* 14: 1929-1945.
- Mantelatto, F.L.M. & Dias, L.L. 1999. Extension of the known distribution of *Charybdis hellerii* (A. Milne-Edwards, 1867) (Decapoda, Portunidae) along the western tropical South Atlantic. *Crustaceana* 72: 617-620.
- Minchin, D., S. Gollasch, A.N. Cohen, C.L. Hewitt & S. Olenin. 2009. Characterizing vectors of marine invasion, p. 109-116. *In* G. Rilov, J.A. Crooks & A. Jeffrey (eds.). Biological invasions in marine ecosystems ecological, management, and geographic

- perspectives. Springer-Verlag, Berlin, Brandenburg, Germany.
- Molnar, J.L., R.L. Gamboa, C. Revenga & M.D. Spalding. 2008. Assessing the global threat of invasive species to marine biodiversity. *Front Ecol. Environ.* 6: 485-492.
- Negreiros-Françoço, M.L. 1996. The zoea I of *Charybdis hellerii* (A. Milne-Edwards, 1867) (Decapoda, Portunidae) obtained in laboratory. *Nauplius* 4: 165-168.
- Oliveira, A.E.S. & J.C. Creed. 2008. Mollusca, Bivalvia, *Isognomon bicolor* (C. B. Adams 1845): Distribution extension. *Check List* 4: 386-388.
- Overstreet, R.M., D.V. Lightner, K. Hasson, S. McIlwain & J.M. Lotz. 1997. Susceptibility to Taura Syndrome Virus of some penaeid shrimp species native to the Gulf of México and the Southeastern United States. *J. Invert. Pathology* 69: 165-176.
- Pantoja, C.R. & D.V. Lightner. 2008. Enfermedades virales, p. 55-116. In V.Q. Morales & J. Cuéllar-Anjel (eds.). *Guía técnica: Patología e inmunología de camarones pendidos*. Programa CYTED Red II-D Vannamei, Panama City, Panama.
- Parker, I.M., D. Simberloff, W.M. Lonsdale, K. Goodell, M. Wonham, P.M. Kareiva, M.H. Williamson, B. Von Holle, P.B. Moyle, J.E. Byers & L. Goldwasser. 1999. Impact: Toward a framework for understanding the ecological effects of invaders. *Biol. Invas.* 1: 3-19.
- Reigada, A.L.D., B.S. Sant'anna, C.M. Zangrande & R.C. Costa. 2006. Macrocrustaceans of non-consolidated sublittoral of the São Vicente Estuarine Bay Complex, São Paulo state, Brazil. *Check List* 2: 84-88.
- Rocha, F.M. 2002. Recrutamento e sucessão de uma comunidade bentônica de mesolitoral dominada pela espécie invasora *Isognomon bicolor* (Bivalvia: Isognomidae) C.B. Adams, 1748 em dois costões rochosos submetidos a diferentes condições de batimento de ondas. MSc. Thesis, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Rio de Janeiro, Brazil.
- Rocha, O., E.L.G. Espíndola, N. Fenerich-Verani, J.R. Verani & A.C. Rietzler. 2005. Espécies invasoras em águas doces - estudos de caso e propostas de manejo. Ed. UFSCar, São Carlos, São Paulo, Brazil.
- Rodrigues, G. & H. Suárez. 2001. Anthropogenic dispersal of decapod crustaceans in aquatic environments. *Interciencia* 26: 282-288.
- Rylov, G. & J.A. Crooks. 2009. Marine bioinvasions: Conservation hazards and vehicles for ecological understanding, p. 3-11. In G. Rilov, J.A. Crooks & A. Jeffrey (eds.). *Biological invasions in marine ecosystems: Ecological, management, and geographic perspectives*. Springer-Verlag Berlin, Brandenburg, Germany.
- Sakai, A.K., F.W. Allendorf, J.S. Holt, D.M. Lodge, J. Molofsky, K.A. With, S. Baughman, R.J. Cabin, J.E. Cohen, N.C. Ellstrand, D.E. McCauley, P. O'Neil, I.M. Parker, J.N. Thompson & S.G. Weller. 2001. The population biology of invasive species. *Ann. Rev. Ecol. Sys.* 32: 305-332.
- Sala, O.E., F.S. Chapin III, J.J. Armesto, R. Berlow, J. Bloomfield, R. Dirzo, E. Huber-Sanwald, L.F. Huenneke, R.B. Jackson, A. Kinzig, R. Leemans, D. Lodge, H.A. Mooney, M. Oesterheld, N.L. Poff, M.T. Sykes, B.H. Walker, M. Walker & D.H. Wall. 2000. Global biodiversity scenarios for the year 2100. *Science* 287: 1770-1774.
- Santos, M.C.F. & P.A. Coelho. 2002. Espécies exóticas de camarões Peneideos (*Penaeus monodon* Fabricius, 1798 e *Litopenaeus vannamei* Boone, 1931) nos ambientes estuarino e marinho do nordeste do Brasil. *Bol. Técn. Cient. CEPENE* 10: 209-222.
- Stein, B., L.S. Kutner & J.S. Adams. 2000. *Precious Heritage: The status of biodiversity in the United States*. Oxford University, Oxfordshire, United Kingdom.
- Tavares, M. & J.B. Mendonça Jr. 1996. *Charybdis hellerii* (A. Milne Edwards, 1867) (Brachyura: Portunidae), eight nonindigenous marine decapod recorded from Brazil. *Crustacean Res.* 25: 151-57.
- Tavares, M. & J.B. Mendonça Jr. 2004. Introdução de crustáceo decápodes exóticos no Brasil: uma roleta ecológica, p. 59-76. In J.S.V. Silva & R.C.C.L. Souza (eds.). *Água de lastro e bioinvasão*. Interciência, Rio de Janeiro, Brazil.
- Valenti, W.C. & M.B. New. 2000. Grow-out systems Monoculture, p. 157-176. In M.B. New & W.C. Valenti (eds.). *Freshwater prawn culture: The farming of *Macrobrachium rosenbergii**. Blackwell Science, Oxfordshire, United Kingdom.
- Wilcove, D.S., D. Rothstein, J. Dubow, A. Phillips & E. Losos. 1998. Quantifying threats to imperiled species in the United States. *BioScience* 48: 607-615.
- Williamson, M. 1996. *Biological invasions*. Chapman & Hall, New York, USA.
- Wilson, E.O. 1992. *The diversity of life*. W.W. Norton, New York, USA.